



Syddansk Universitet

Charges in POPC Monolayers - a Temperature Dependant Monolayer Study

Hermanowska, Malgorzata; Bijelic, Goran; Claesson, Per; Klösger, Beate

Published in:
Biophysical Journal

Publication date:
2008

Document Version
Final published version

[Link to publication](#)

Citation for pulished version (APA):

Hermanowska, M., Bijelic, G., Claesson, P., & Klösger, B. (2008). Charges in POPC Monolayers - a Temperature Dependant Monolayer Study. *Biophysical Journal*, (2).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Joint Meeting of the Biophysical Society, 52nd Annual Meeting and 16th International Biophysical Congress (IUPAB),
February 2-6, 2008,
Long Beach, California, USA.*

Charges in POPC Monolayers - a Temperature Dependant Monolayer Study

Malgorzata Hermanowska¹, Goran Bijelic², Beate Klösgen¹, Per Claesson².

¹University of Southern Denmark, Odense, Denmark, ²Royal Institute of Technology, Stockholm, Sweden.

Abstract: The surface of a bio-membrane is the first site of interaction for many biophysico-chemical processes. The access to the surface, and the local structural modifications induced to the lipid bilayer upon interaction with an adsorbing guest molecule must be influenced significantly by the presence of charges, and local changes in surface charge density. Here we present preliminary results from an isothermal compression study of POPC monolayers into which charges were inoculated by addition of the ethylated lipid variety. The phase state will be discussed in view of its relevance surface adsorption.

The phospholipids examined in this study are: 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-sn-glycero-3-ethylphosphocholine (EPOPC). Charge effects on POPC monolayers at the air-water interface were investigated by addition of different percentages of the cationic EPOPC. These two phospholipids have the same molecular sequence apart from a slight difference in the headgroup region. The isotherms obtained for the mixtures are compared with pure POPC monolayer isotherms as a standard system.

Temperature dependence of the isotherms for both pure POPC and the different monolayer compositions were examined at temperatures ranging from 5°C to 50°C. The π -A isotherms for POPC exposed a behaviour which agrees well with established theoretical approaches for temperatures above 20°C: in that temperature region the surface pressure is increased as temperature is raised. However at temperatures lower than 20°C non-ideal behaviour is observed in that surface pressure increases again upon decrease in temperature. The minimum in surface pressure as a function of temperature is seen around 20°C. The increase in monolayer's surface pressure upon introduction of a surface charge indicates repulsive interaction between molecules, this might decrease stability of film thus, induce curvature frustration and might destabilize a bilayer formed from such a mixture.